



Attorney's Docket: 1999DE132
 Serial No.: 09/722,760
 Art Unit: 1753

4) (Original) The method as claimed in claim 1, wherein the low molecular weight organic cation is a substituted ammonium, phosphonium, thionium or triphenylcarbonium ion or a cationic metal complex.


5) (Previously Amended) The method as claimed in claim 4, wherein the ammonium ion has one of the formulae (a) - (j)




(a)




(b)




(c)




(d)



(e)




(f)




(g)


Attorney's Docket: 1999DE132
 Serial No.: 09/722,760
 Art Unit: 1753



(h)



(i)




(j)

in which

R^1 to R^{18} are identical or different and represent hydrogen, CN, $(\text{CH}_2)_{1-18}\text{CN}$, halogen, branched or unbranched C_1 - C_{32} -alkyl, mono- or polyunsaturated C_2 - C_{32} -alkenyl, C_1 - C_{22} -alkoxy, C_1 - C_{22} -hydroxyalkyl, C_1 - C_{22} -halogenoalkyl, C_2 - C_{22} -halogenoalkenyl, C_1 - C_{22} -aminoalkyl, $(\text{C}_1$ - C_{12})-trialkyl-ammonium- $(\text{C}_1$ - C_{22})-alkyl; $(\text{C}_1$ - C_{22})-alkylene- $(\text{C}=\text{O})\text{O}$ - $(\text{C}_1$ - C_{32})-alkyl, $(\text{C}_1$ - C_{22})-alkylene- $(\text{C}=\text{O})\text{O}$ -aryl, $(\text{C}_1$ - C_{22})-alkylene- $(\text{C}=\text{O})\text{NH}$ - $(\text{C}_1$ - C_{32})-alkyl, $(\text{C}_1$ - C_{22})-alkylene- $(\text{C}=\text{O})\text{NH}$ -aryl, $(\text{C}_1$ - C_{22})-alkylene- $\text{O}(\text{CO})$ - $(\text{C}_1$ - C_{32})-alkyl, $(\text{C}_1$ - C_{22})-alkylene- $\text{O}(\text{CO})$ -aryl, $(\text{C}_1$ - C_{22})-alkylene- $\text{NH}(\text{C}=\text{O})$ - $(\text{C}_1$ - C_{32})-alkyl, $(\text{C}_1$ - C_{22})-alkylene- NHCO -aryl,

wherein



are optionally inserted into the acid ester or acid amide bonds;

$[(\text{C}_1$ - $\text{C}_{12})$ -alkylene- $\text{O}]_{1-100}$ -H; aryl, $(\text{C}_1$ - $\text{C}_{18})$ -alkylenearyl; $-(\text{O}-\text{SiR}'_2)_{1-32}-\text{O}-\text{SiR}'_3$, in which R' has the meaning C_1 - C_{12} -alkyl, phenyl, benzyl or C_1 - C_{12} -alkoxy; heterocyclyl, C_1 - C_{18} -alkylene-heterocyclyl, wherein the aryl and heterocyclyl radicals are optionally mono- or polysubstituted on carbon atoms or heteroatoms by C_1 - C_{12} -alkyl, C_1 - C_4 -alkenyl, C_1 - C_4 -alkoxy, hydroxy- $(\text{C}_1$ - C_4)-alkyl, amino- $(\text{C}_1$ - C_4)-alkyl, C_1 - C_4 -alkylimino,

Attorney's Docket: 1999DE132


Serial No.: 09/722,760

Art Unit: 1753

carboxyl, hydroxyl, amino, nitro, cyano, halogen, C₁-C₁₂-acyl, C₁-C₄-halogenoalkyl, C₁-C₄-alkylcarbonyl, C₁-C₄-alkylcarbonyloxy, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkylaminocarbonyl, C₁-C₄-alkylcarbonylimino, C₆-C₁₀-arylcarbonyl, aminocarbonyl, aminosulfonyl, C₁-C₄-alkylaminosulfonyl, phenyl, naphthyl, or heteroaryl[.];

R¹⁹ represents C₄-C₁₁-alkylene, -(C₂H₄-O-)₁₋₁₇-(CH₂)₁₋₂-, -(C₂H₄-NR-)₁₋₁₇-(CH₂)₁₋₂-, in which R is hydrogen or C₁-C₁₂-alkyl;

X has the meaning of Y or -CO-CH₂-CO-,





Y has the meaning $\begin{array}{c} \text{C} \\ \parallel \\ \text{O} \end{array}$, $\begin{array}{c} \text{C} \\ \parallel \\ \text{S} \end{array}$, $\begin{array}{c} \text{C} \\ \parallel \\ \text{NH} \end{array}$, -(CH₂)₁₋₁₈-,

Attorney's Docket: 1999DE132

Serial No.: 09/722,760

Art Unit: 1753



or o-, p-, m-(C₆-C₁₄)-arylene or (C₄-C₁₄)-heteroarylene with 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, S and a combination thereof;

R⁶⁰ represents C₁-C₃₂-acyl, C₁-C₂₂-alkyl, C₂-C₂₂-alkenyl, C₁-C₁₈-alkylene-C₆-C₁₀-aryl, C₁-C₂₂-alkylene-heterocyclyl, C₆-C₁₀-aryl or (C₄-C₁₄)-heteroaryl with 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, S, and a combination thereof;


R⁶¹ and R⁶⁴ represent -(CH₂)₁₋₁₈-, C₁-C₁₂-alkylene-C₆-C₁₀-arylene, C₆-C₁₀-arylene, C₀-C₁₂-alkylene-heterocyclyl;

Z represents -NH- or -O-;

A₁⁻ and A₃⁻ represent -COO⁻, -SO₃⁻, -OSO₃⁻, -SO₂⁻, -COS⁻ or -CS₂⁻;

A₂ represents -SO₂Na, -SO₃Na, -SO₂H, -SO₃H or hydrogen;

R⁶⁹ and R⁷⁰ independently of one another represent hydrogen, C₁-C₃₂-alkyl, in which the alkyl chain optionally contain one or more of the groups -NH-CO-, -CO-NH-, -CO-O- or -O-CO-; C₁-C₁₈-alkylene-aryl, C₀-C₁₈-alkylene-heterocyclyl, C₁-C₁₈-hydroxyalkyl, C₁-C₁₈-halogenoalkyl, aryl, -(CH₂)₃-SO₃⁻;



R⁷¹ and R⁷² represent -(CH₂)₁₋₁₂-; and

Attorney's Docket: 1999DE132
Serial No.: 09/722,760
Art Unit: 1753

R⁷³ and R⁷⁴ represent hydrogen or C₁-C₂₂-alkyl.

6) (Original) The method as claimed in claim 5, wherein R¹ to R¹⁸ denote hydrogen CN, CH₂-CN, CF₃, C₁-C₂₂-alkyl, C₂-C₁₈-alkenyl, C₁-C₁₈-alkoxy, C₁-C₁₈-hydroxy-alkyl, C₁-C₁₈-halogenoalkyl, C₂-C₁₈-halogenoalkenyl, C₁-C₁₈-aminoalkyl, (C₁-C₈)-trialkylammonium-(C₁-C₁₈)-alkyl, (C₁-C₁₈)-alkylene-O(C=O)-(C₁-C₂₂)alkyl, (C₁-C₁₈)-alkylene-O(C=O)-phenyl, (C₁-C₁₈)-alkylene-NHCO-(C₁-C₂₂)alkyl, (C₁-C₁₈)-alkylene-NHCO-phenyl, (C₁-C₁₈)-alkylene-(C=O)O-(C₁-C₂₂)alkyl, (C₁-C₁₈)-alkylene-(C=O)O-phenyl, (C₁-C₁₈)-alkylene-(C=O)NH-(C₁-C₂₂)alkyl, (C₁-C₁₈)-alkylene-CONH-phenyl, benzyl, phenyl, naphthyl, C₁-C₁₂-alkylene-heterocyclyl;

R¹⁹ denotes C₄-C₈-alkylene, -(C₂H₄-O)₁₋₉-(CH₂)₁₋₂ or -(C₂H₄-NH)₁₋₉-(CH₂)₁₋₂;

R⁶⁰ denotes C₁-C₁₈-acyl, C₁-C₁₈-alkyl, C₂-C₁₈-alkenyl, C₁-C₁₂-alkylene-phenyl, C₁-C₁₈-alkylene-pyridyl, phenyl or pyridyl;

R⁶¹ and R⁶⁴ denote -(CH₂)₁₋₁₂, C₁-C₈-alkylene-phenylene, phenylene or C₁-C₈-alkylenepyridylene or piperidylene;

R⁷¹ and R⁷² denote -(CH₂)₁₋₈ and

R⁷³ and R⁷⁴ denote hydrogen or (C₁-C₁₈)-alkyl.

7) (Previously Amended) The method as claimed in claim 4, wherein the ammonium ion is an aliphatic or aromatic 5- to 12-membered heterocyclic radical with 1 to 4 atoms selected from the group consisting of N, O and S, or a combination thereof, belonging to the rings.


8) (Original) The method as claimed in claim 7, wherein the heterocyclic radical is pyridinium, pyridazinium, pyrimidinum, pyrazinium, purinium, tetraazaporphyrinium, piperidinium, morpholinum, tetrazonum, triaza-cyclononanium or tetraaza-cyclododecanum.

Attorney's Docket: 1999DE132
 Serial No.: 09/722,760
 Art Unit: 1753

9) (Original) The method as claimed in claim 4, wherein the cationic metal complex is a metal carboxylate, metal salicylate, metal sulfonate, 1:1 metal-azo complex or a metal dithiocarbamate.

10) (Previously Amended) The method as claimed in claim 9, wherein the metal is selected from the group consisting of Al, Mg, Ca, Sr, Ba, TiO, VO, Cr, V, Ti, Zr, Sc, Mn, Fe, Co, Ni, Cu, Zn and ZrO.

11) (Original) The method as claimed in claim 1, wherein the organic cation is a fluorinated ammonium ion of the formula (x)



in which

R^{28} denotes perfluorinated alkyl having 5 to 11 carbon atoms and

R^{29} , R^{30} and R^{31} are identical or different and denote alkyl having 1 to 5 carbon atoms.

12) (Original) Salt-like structured silicate, in which the silicate is hectorite, beldellite, illite, muscovite, xantophyllite, margarite, sepiolite, saponite, mica, feldspar, nontronite, montmorillonite, smectite, bentonite, faujasite, zeolite A, X or Y, permutite, sasil or a combination thereof; and the cation is an ion of the formula (x) as claimed in claim 9.

Attorney's Docket: 1999DE132
Serial No.: 09/722,760
Art Unit: 1753

13) (Original) A process for the preparation of a salt-like structured silicate as claimed in claim 12, which comprises combining the silicate and a salt of the cation of formula (x) in an aqueous medium.

14) (Previously Amended) An electrophotographic toner comprising 30 to 99.99% by weight of a binder, and 0.01 to 50% by weight, of at least one salt of ionic structured silicates in which the cation is a low molecular weight organic cation and the anion is an island, cyclic, group, chain, ribbon, laminar or matrix silicate or a combination thereof, based on the total weight of the electrophotographic toner.

15) (Previously Amended) An electrophotographic toner as claimed in claim 14, comprising 40 to 99.5% by weight of a binder, and 0.05 to 20% by weight of at least one salt of ionic structured silicates in which the cation is a low molecular weight organic cation and the anion is an island, cyclic, group, chain, ribbon, laminar or matrix silicate or a combination thereof, based on the total weight of the electrophotographic toner.

16) (Previously Added) The method of claim 4, wherein the ammonium ion is an aliphatic or aromatic 5- to 12-membered heterocyclic radical with 1 to 4 atoms selected from the group consisting of N, O and S, or a combination thereof, belonging to the rings, wherein 2 to 8 rings are fused.

17) (Previously Added) The method as claimed in claim 9, wherein the metal is selected from the group consisting of Al, Mg, Ca, Sr, Ba, TiO, VO, Cr, V, Ti, Zr, Sc, Mn, Fe, Co, Ni, Cu, Zn and ZrO, and the metal complex contains one or more further ligands.

C¹
18) (Currently Amended) A method of imparting, controlling or improving the charge of an electrophotographic toner or developer, of a powder coating, or of an electret material, comprising the steps of adding a salt structured silicate in which the cation

Attorney's Docket: 1999DE132
Serial No.: 09/722,760
Art Unit: 1753

is NH_4^+ , H_3O^+ , an alkali metal, alkaline earth metal, earth metal or transition metal ion or a low molecular weight organic cation or a combination thereof and the anion is an island, cyclic, group, chain, ribbon, laminar or matrix silicate or a combination thereof to a binder of an electrophotographic toner or developer or of a powder coating, or to an electret material.

19. (Previously Amended) An electrophotographic toner or developer comprising distearyldimethyl ammonium bentonite.
20. (Previously Added) The electrophotographic toner as claimed in claim 14, further comprising 0.001 to 50% by weight, of a coloring agent, based on the total weight of the electrophotographic toner.
21. (Previously Amended) A composition comprising 30 to 99.99% by weight of a binder, and 0.01 to 50% by weight, of at least one salt of ionic structured silicates in which the cation is a low molecular weight organic cation and the anion is an island, cyclic, group, chain, ribbon, laminar or matrix silicate or a combination thereof, based on the total weight of the composition, wherein the composition is an electrophotographic toner.
22. (Previously Added) A method of imparting, controlling or improving the charge of an electrophotographic toner or developer, or an electret material comprising the step of adding a distearyldimethyl ammonium bentonite to a binder of an electrophotographic toner or developer or of a powder coating or of an electret material.